

# References: Introduction to Integrated Behavioral Health Services

Patrick Van Wyk, Ph.D.

1. Katon, W., Von Korff, M., Lin, E., Walker, E., Simon, G. E., Bush, T., ... & Russo, J. (1995). Collaborative management to achieve treatment guidelines: impact on depression in primary care. *Jama*, 273(13), 1026-1031.
2. Kroenke, K., & Mangelsdorff, A. D. (1989). Common symptoms in ambulatory care: incidence, evaluation, therapy, and outcome. *The American journal of medicine*, 86(3), 262-266.
3. Narrow, W. E., Regier, D. A., Rae, D. S., Manderscheid, R. W., & Locke, B. Z. (1993). Use of services by persons with mental and addictive disorders: Findings from the National Institute of Mental Health Epidemiologic Catchment Area Program. *Archives of General Psychiatry*, 50(2), 95.
4. Kessler, R. C., Demler, O., Frank, R. G., Olfson, M., Pincus, H. A., Walters, E. E., ... & Zaslavsky, A. M. (2005). Prevalence and treatment of mental disorders, 1990 to 2003. *New England Journal of Medicine*, 352(24), 2515-2523.
5. Kessler, R. C., Demler, O., Frank, R. G., Olfson, M., Pincus, H. A., Walters, E. E., ... & Zaslavsky, A. M. (2005). Prevalence and treatment of mental disorders, 1990 to 2003. *New England Journal of Medicine*, 352(24), 2515-2523.
6. Fisher, L., & Ransom, D. C. (1996). Developing a strategy for managing behavioral health care within the context of primary care. *Archives of family medicine*, 6(4), 324-333.
7. Hoge, C. W., Auchterlonie, J. L., & Milliken, C. S. (2006). Mental health problems, use of mental health services, and attrition from military service after returning from deployment to Iraq or Afghanistan. *Jama*, 295(9), 1023-1032.
8. Cunningham, P. J. (2009). Beyond parity: primary care physicians' perspectives on access to mental health care. *Health Affairs*, 28(3), w490-w501.
9. JOINT PRINCIPLES: INTEGRATING BEHAVIORAL HEALTH CARE INTO THE PATIENT-CENTERED MEDICAL HOME *Annals of Family Medicine*, 2014(12) 183-185

10. Peek, C. J. (2011). A collaborative care lexicon for asking practice and research development questions. In C. Mullican (Ed.), *AHRQ Publication No. 11-0067* (pp. 25-44). Rockville, MD: Agency for Healthcare Research and Quality.
11. Chen, H., Coakley, E. H., Cheal, K., Maxwell, J., Costantino, G., Krahn, D. D., ... & Levkoff, S. E. (2006). Satisfaction With Mental Health Services in Older Primary Care Patients. *American Journal of Geriatric Psych*, 14(4), 371-379.
12. Bartels, S. J., Coakley, E. H., Zubritsky, C., Ware, J. H., Miles, K. M., Areán, P. A., ... & Levkoff, S. E. (2004). Improving access to geriatric mental health services: a randomized trial comparing treatment engagement with integrated versus enhanced referral care for depression, anxiety, and at-risk alcohol use. *American Journal of Psychiatry*, 161(8), 1455-1462.
13. Kwan, B.M. & Nease, D.E. (2013). The state of the evidence for integrated behavioral health in primary care. In: M.R. Talen & A .Burke Valeras (Eds.), *Integrated Behavioral Health in Primary Care: Evaluating the Evidence, Identifying the Essentials* (pp. 65-98). New York: Springer.
14. Gilbody, S., Bower, P., Fletcher, J., Richards, D., & Sutton, A. J. (2006). Collaborative Care for DepressionA Cumulative Meta-analysis and Review of Longer-term Outcomes. *Archives of Internal Medicine*, 166(21), 2314-2321.
15. Gulliksson, M., Burell, G., Vessby, B., Lundin, L., Toss, H., & Svärdsudd, K. (2011). Randomized controlled trial of cognitive behavioral therapy vs standard treatment to prevent recurrent cardiovascular events in patients with coronary heart disease: Secondary Prevention in Uppsala Primary Health Care project (SUPRIM). *Archives of Internal Medicine*, 171(2), 134-140.
16. Schweikert, B., E. Jacobi, et al. (2006) Effectiveness and cost-effectiveness of adding a cognitive behavioral treatment to the rehabilitation of chronic low back pain (Structured abstract). *Journal of Rheumatology Volume*, 2519-2526
17. Roux, L., K. M. Kuntz, et al. (2006) Economic evaluation of weight loss interventions in overweight and obese women (Structured abstract). *Obesity Volume*, 1093-1106
18. Butler M, Kane RL, McAlpine D, Kathol, RG, Fu SS, Hagedorn H, Wilt TJ. Integration of Mental Health/Substance Abuse and Primary Care No. 173 AHRQ Publication No. 09-E003. Rockville, MD. Agency for Healthcare Research and Quality. October 2008.
19. Blount, F. A., & Miller, B. F. (2009). Addressing the workforce crisis in integrated primary care. *Journal of Clinical Psychology in Medical Settings*, 16(1), 113-119.

20. Kessler, R., & Stafford, D. (2008). Primary care is the de facto mental health system. In *Collaborative medicine case studies* (pp. 9-21). Springer New York.
21. Strosahl, K., & Robinson, P. (2008). The primary care behavioral health model: Applications to prevention, acute care and chronic condition management. In *Collaborative medicine case studies* (pp. 85-95). Springer New York.
22. Unutzer, J., W. J. Katon, et al. (2008) Long-term cost effects of collaborative care for late-life depression (Brief record). *American Journal of Managed Care Volume*, 95-100
23. Lindgren, P., J. Lindstrom, et al. (2007) Lifestyle intervention to prevent diabetes in men and women with impaired glucose tolerance is cost-effective (Structured abstract). *International Journal of Technology Assessment in Health Care Volume*, 177-183
24. Sommers, L. S., Marton, K. I., Barbaccia, J. C., & Randolph, J. (2000). Physician, nurse, and social worker collaboration in primary care for chronically ill seniors. *Archives of Internal medicine*, 160(12), 1825-1833.
25. Blumenthal, J. A., Babyak, M., Wei, J., O'Connor, C., Waugh, R., Eisenstein, E., ... & Reed, G. (2002). Usefulness of psychosocial treatment of mental stress-induced myocardial ischemia in men. *The American journal of cardiology*, 89(2), 164-168.
26. Ciechanowski, P. S., Katon, W. J., & Russo, J. E. (2000). Depression and diabetes: impact of depressive symptoms on adherence, function, and costs. *Archives of internal medicine*, 160(21), 3278-3285.
27. Peek, C. J. National Integration Academy Council.(2013). Lexicon for behavioral health and primary care integration: Concepts and definitions developed by expert consensus (AHRQ Publication No. 13-IP001-EF).
28. Ratanawongsa, N., Roter, D., Beach, M. C., Laird, S. L., Larson, S. M., Carson, K. A., & Cooper, L. A. (2008). Physician burnout and patient-physician communication during primary care encounters. *Journal of general internal medicine*, 23(10), 1581-1588.
29. Vahey, D. C., Aiken, L. H., Sloane, D. M., Clarke, S. P., & Vargas, D. (2004). Nurse burnout and patient satisfaction. *Medical care*, 42(2 Suppl), II57.
30. Mosen, D. M., Schmittdiel, J., Hibbard, J., Sobel, D., Remmers, C., & Bellows, J. (2007). Is patient activation associated with outcomes of care for adults with chronic conditions?. *The Journal of ambulatory care management*, 30(1), 21-29.
31. Cleland, J. G., & Ekman, I. (2010). Enlisting the help of the largest health care workforce—patients. *JAMA*, 304(12), 1383-1384.

32. Hibbard, J. 2008. Increasing Patient Activation to Improve Health and Reduce Costs. Institute for Policy Research and Innovation, University of Oregon
33. Center for studying health system change 2007
34. Barnes KA, Kroening-Roche, JC, Comfort BW (2012). The developing vision of primary care. *The New England Journal of Medicine*, 367, 891-893.
35. McGinnis JM, Williams-Russo P, Knickman JR. (2002). The case for more active policy attention to health promotion. *Health Affairs*, 21, 78-93.
36. Mokdad, A. H., Marks, J. S., Stroup, D. F., & Gerberding, J. L. (2004). Actual causes of death in the United States, 2000. *Jama*, 291(10), 1238-1245.
- 36.1. Kathol, R. G., Melek, S., Bair, B., & Sargent, S. (2008). Financing mental health and substance use disorder care within physical health: a look to the future. *Psychiatric Clinics of North America*, 31(1), 11-25.
37. Manderscheid, R., & Kathol, R. (2014). Fostering sustainable, integrated medical and behavioral health services in medical settings. *Annals of Internal Medicine*, 160, 61-65.
38. Kathol, R. G., & Rollman, B. L. (2014). Value-based financially sustainable behavioral health components in patient-centered medical homes. *The Annals of Family Medicine*, 12(2), 172-175.
39. Fisher, L., & Dickinson, W. P. (2014). Psychology and primary care: New collaborations for providing effective care for adults with chronic health conditions. *American Psychologist*, 69(4), 355.
40. Blount, A., Schoenbaum, M., Kathol, R., Rollman, B. L., Thomas, M., O'Donohue, W., & Peek, C. J. (2007). The economics of behavioral health services in medical settings: A summary of the evidence. *Professional Psychology: Research and Practice*, 38(3), 290.
41. Kessler, R. (2008). Integration of care is about money too: The health and behavior codes as an element of a new financial paradigm. *Families, Systems, & Health*, 26(2), 207.